

# Background knowledge

- Contents:**
- A** Operations with surds (radicals)
  - B** Standard form (scientific notation)
  - C** Number systems and set notation
  - D** Algebraic simplification
  - E** Linear equations and inequalities
  - F** Absolute value (modulus)
  - G** Product expansion
  - H** Factorisation
    - Investigation 1: Another factorisation technique*
  - I** Formula rearrangement
  - J** Adding and subtracting algebraic fractions
  - K** Congruence and similarity
  - L** Coordinate geometry
    - Investigation 2: Finding where lines meet using technology*



This section contains material that is normally covered prior to this course. It is assumed, background knowledge. Not all preliminaries are covered within it. However, other necessary work is revised within the chapters which follow this one.

## A OPERATIONS WITH SURDS (RADICALS)

Real numbers like  $\sqrt{2}$ ,  $\sqrt{3}$ ,  $\sqrt{5}$ ,  $\sqrt{6}$ , etc., are called **surds** or **radicals**. Surds are present in solutions to some quadratic equations.  $\sqrt{4}$  is not a surd as it simplifies to 2. Surds are irrational real numbers.

**Definition:**  $\sqrt{a}$  is the non-negative number such that  $\sqrt{a} \times \sqrt{a} = a$ .

**Properties:**

- $\sqrt{a}$  is never negative, that is,  $\sqrt{a} \geq 0$ .
- $\sqrt{a}$  is meaningful only for  $a \geq 0$ .
- $\sqrt{ab} = \sqrt{a} \times \sqrt{b}$  for  $a \geq 0$  and  $b \geq 0$ .
- $\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$  for  $a \geq 0$  and  $b > 0$ .

### SURDIC OPERATIONS

#### Example 1

Write as a single surd:      **a**  $\sqrt{2} \times \sqrt{3}$                       **b**  $\frac{\sqrt{18}}{\sqrt{6}}$

<b>a</b> $\sqrt{2} \times \sqrt{3}$ $= \sqrt{2 \times 3}$ $= \sqrt{6}$	<b>b</b> $\frac{\sqrt{18}}{\sqrt{6}}$ $= \sqrt{\frac{18}{6}}$ $= \sqrt{3}$	<b>or</b> $\frac{\sqrt{18}}{\sqrt{6}}$ $= \frac{\sqrt{6 \times 3}}{\sqrt{6}}$ $= \sqrt{3}$
--	--	--

### EXERCISE A

1 Write as a single surd or rational number:

<b>a</b> $\sqrt{3} \times \sqrt{5}$	<b>b</b> $(\sqrt{3})^2$	<b>c</b> $2\sqrt{2} \times \sqrt{2}$	<b>d</b> $3\sqrt{2} \times 2\sqrt{2}$
<b>e</b> $3\sqrt{7} \times 2\sqrt{7}$	<b>f</b> $\frac{\sqrt{12}}{\sqrt{2}}$	<b>g</b> $\frac{\sqrt{12}}{\sqrt{6}}$	<b>h</b> $\frac{\sqrt{18}}{\sqrt{3}}$

#### Example 2

Simplify:      **a**  $3\sqrt{3} + 5\sqrt{3}$                       **b**  $2\sqrt{2} - 5\sqrt{2}$

<b>a</b> $3\sqrt{3} + 5\sqrt{3}$ $= (3 + 5)\sqrt{3}$ $= 8\sqrt{3}$	<b>b</b> $2\sqrt{2} - 5\sqrt{2}$ $= (2 - 5)\sqrt{2}$ $= -3\sqrt{2}$
--	---

Compare with  
 $2x - 5x = -3x$



2 Simplify the following mentally:

**a**  $2\sqrt{2} + 3\sqrt{2}$

**b**  $2\sqrt{2} - 3\sqrt{2}$

**c**  $5\sqrt{5} - 3\sqrt{5}$

**d**  $5\sqrt{5} + 3\sqrt{5}$

**e**  $3\sqrt{5} - 5\sqrt{5}$

**f**  $7\sqrt{3} + 2\sqrt{3}$

**g**  $9\sqrt{6} - 12\sqrt{6}$

**h**  $\sqrt{2} + \sqrt{2} + \sqrt{2}$

### Example 3

Write  $\sqrt{18}$  in the form  $a\sqrt{b}$  where  $a$  and  $b$  are integers,  $a$  is as large as possible.

$$\begin{aligned}\sqrt{18} &= \sqrt{9 \times 2} && \{9 \text{ is the largest perfect square factor of } 18\} \\ &= \sqrt{9} \times \sqrt{2} \\ &= 3\sqrt{2}\end{aligned}$$

3 Write the following in the form  $a\sqrt{b}$  where  $a$  and  $b$  are integers and  $a$  is as large as possible:

**a**  $\sqrt{8}$

**b**  $\sqrt{12}$

**c**  $\sqrt{20}$

**d**  $\sqrt{32}$

**e**  $\sqrt{27}$

**f**  $\sqrt{45}$

**g**  $\sqrt{48}$

**h**  $\sqrt{54}$

**i**  $\sqrt{50}$

**j**  $\sqrt{80}$

**k**  $\sqrt{96}$

**l**  $\sqrt{108}$

### Example 4

Simplify:  $2\sqrt{75} - 5\sqrt{27}$

$$\begin{aligned}2\sqrt{75} - 5\sqrt{27} &= 2\sqrt{25 \times 3} - 5\sqrt{9 \times 3} \\ &= 2 \times 5 \times \sqrt{3} - 5 \times 3 \times \sqrt{3} \\ &= 10\sqrt{3} - 15\sqrt{3} \\ &= -5\sqrt{3}\end{aligned}$$

4 Simplify:

**a**  $4\sqrt{3} - \sqrt{12}$

**b**  $3\sqrt{2} + \sqrt{50}$

**c**  $3\sqrt{6} + \sqrt{24}$

**d**  $2\sqrt{27} + 2\sqrt{12}$

**e**  $\sqrt{75} - \sqrt{12}$

**f**  $\sqrt{2} + \sqrt{8} - \sqrt{32}$

### Example 5

Write  $\frac{9}{\sqrt{3}}$  without a radical in the denominator.

$$\begin{aligned}\frac{9}{\sqrt{3}} &= \frac{9}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} \\ &= \frac{9\sqrt{3}}{3} \\ &= 3\sqrt{3}\end{aligned}$$

5 Write without a radical in the denominator:

a  $\frac{1}{\sqrt{2}}$

b  $\frac{6}{\sqrt{3}}$

c  $\frac{7}{\sqrt{2}}$

d  $\frac{10}{\sqrt{5}}$

e  $\frac{10}{\sqrt{2}}$

f  $\frac{18}{\sqrt{6}}$

g  $\frac{12}{\sqrt{3}}$

h  $\frac{5}{\sqrt{7}}$

i  $\frac{14}{\sqrt{7}}$

j  $\frac{2\sqrt{3}}{\sqrt{2}}$

## B STANDARD FORM (SCIENTIFIC NOTATION)

**Standard form** (or **scientific notation**) involves writing any given number as a number between 1 and 10, multiplied by a power of 10,

i.e.,  $a \times 10^n$  where  $a$  lies between 1 and 10, i.e.,  $1 \leq a < 10$ .

### Example 6

Write in standard form:      a 37 600      b 0.000 86

a  $37\,600 = 3.76 \times 10\,000$  {shift decimal point 4 places to the left and  $\times 10\,000$ }  
 $= 3.76 \times 10^4$

b  $0.000\,86 = 8.6 \div 10^4$  {shift decimal point 4 places to the right and  $\div 10\,000$ }  
 $= 8.6 \times 10^{-4}$

### EXERCISE B

1 Express the following in standard form (scientific notation):

a 259

b 259 000

c 2.59

d 0.259

e 0.000 259

f 40.7

g 4070

h 0.0407

i 407 000

j 407 000 000

k 0.000 0407

2 Express the following in standard form (scientific notation):


a The distance from the Earth to the Sun is 149 500 000 000 m.

b Bacteria are single cell organisms, some of which have a diameter of 0.000 3 mm.

c A speck of dust is smaller than 0.001 mm.

d The central temperature of the Sun is 15 million degrees Celsius.

e A single red blood cell lives for about four months and during this time it will circulate around the body 300 000 times.



### Example 7

Write as an ordinary number:

a  $3.2 \times 10^2$

b  $5.76 \times 10^{-5}$

a  $3.2 \times 10^2$   
 $= 3.20 \times 100$   
 $= 320$

b  $5.76 \times 10^{-5}$   
 $= 000005.76 \div 10^5$   
 $= 0.000\,057\,6$

3 Write as an ordinary decimal number:

- a**  $4 \times 10^3$       **b**  $5 \times 10^2$       **c**  $2.1 \times 10^3$       **d**  $7.8 \times 10^4$   
**e**  $3.8 \times 10^5$       **f**  $8.6 \times 10^1$       **g**  $4.33 \times 10^7$       **h**  $6 \times 10^7$

4 Write as an ordinary decimal number:

- a**  $4 \times 10^{-3}$       **b**  $5 \times 10^{-2}$       **c**  $2.1 \times 10^{-3}$       **d**  $7.8 \times 10^{-4}$   
**e**  $3.8 \times 10^{-5}$       **f**  $8.6 \times 10^{-1}$       **g**  $4.33 \times 10^{-7}$       **h**  $6 \times 10^{-7}$

5 Write as an ordinary decimal number:

- a** The wave length of light is  $9 \times 10^{-7}$  m.  
**b** The estimated world population for the year 2000 is  $6.130 \times 10^9$ .  
**c** The diameter of our galaxy, the Milky Way, is  $1 \times 10^5$  light years.  
**d** The smallest viruses are  $1 \times 10^{-5}$  mm in size.

6 Find, with decimal part correct to 2 places:

- a**  $(3.42 \times 10^5) \times (4.8 \times 10^4)$       **b**  $(6.42 \times 10^{-2})^2$       **c**  $\frac{3.16 \times 10^{-10}}{6 \times 10^7}$   
**d**  $(9.8 \times 10^{-4}) \div (7.2 \times 10^{-6})$       **e**  $\frac{1}{3.8 \times 10^5}$       **f**  $(1.2 \times 10^3)^3$

7 If a missile travels at 5400 km/h how far will it travel in:

- a** 1 day      **b** 1 week      **c** 2 years?



[Give your answers in standard form with decimal part correct to 2 places and assume that 1 year  $\doteq$  365.25 days.]

8 Light travels at a speed of  $3 \times 10^8$  metres per second. How far will light travel in:

- a** 1 minute      **b** 1 day      **c** 1 year?


[Give your answers with decimal part correct to 2 decimal places and assume that 1 year  $\doteq$  365.25 days.]

## C

## NUMBER SYSTEMS AND SET NOTATION

### NUMBER SYSTEMS

We will use •  $\mathcal{R}$  to represent the set of all **real numbers**. These are all the numbers on the number line.



- $\mathcal{N}$  to represent the set of all **natural numbers**.  $\mathcal{N} = \{0, 1, 2, 3, 4, 5, \dots\}$

- $\mathcal{Z}$  to represent the set of all **integers**.  $\mathcal{Z} = \{0, \pm 1, \pm 2, \pm 3, \pm 4, \dots\}$

**Note:**  $\mathcal{Z}^+$  is the set of all positive integers.  $\mathcal{Z}^+ = \{1, 2, 3, 4, \dots\}$

- $\mathcal{Q}$  to represent the set of all **rational numbers** which are any numbers of the form  $\frac{p}{q}$  where  $p$  and  $q$  are integers,  $q \neq 0$ .

## SET NOTATION

$\{x : -3 < x < 2\}$  reads “the set of all values that  $x$  can be such that  $x$  lies between  $-3$  and  $2$ ”.

Later in the text we will write this as  $x \in ]-3, 2[$ .

*the set of all*      *such that*

## EXERCISE C

1 Write verbal statements for the meaning of:

**a**  $\{x: x > 5\}$

**b**  $\{x: x \leq 3\}$

**c**  $\{y: 0 < y < 6\}$

**d**  $\{x: 2 \leq x \leq 4\}$



**e**  $\{t: 1 < t < 5\}$

**f**  $\{n: n < 2 \text{ or } n \geq 6\}$

**Note:** If a number set like  $\mathbf{N}$ ,  $\mathbf{Z}$  or  $\mathbf{Q}$  is not given we assume we are referring to real numbers (i.e., in  $\mathcal{R}$ ).

## Example 8

Write in set notation:







**a**  $\{x: x \in \mathbf{N}, 1 \leq x \leq 4\}$

**b**  $\{x: -3 \leq x < 4\}$

or  $\{x: x \in \mathbf{Z}, 1 \leq x \leq 4\}$

**Note:**  $\in$  is used to mean “is in”

2 Write in set notation:



3 Sketch the following number sets:

**a**  $\{x: x \in \mathbf{N}, 4 \leq x < 10\}$

**b**  $\{x: x \in \mathbf{Z}, -4 < x \leq 5\}$

**c**  $\{x: x \in \mathcal{R}, -5 < x \leq 4\}$

**d**  $\{x: x \in \mathbf{Z}, x > -4\}$

**e**  $\{x: x \in \mathcal{R}, x \leq 8\}$

## D

## ALGEBRAIC SIMPLIFICATION

Recall that  $a(b + c) = ab + ac$  and  $a(b - c) = ab - ac$ , the **distributive law**.

## EXERCISE D

1 Simplify if possible:

**a**  $3x + 7x - 10$

**b**  $3x + 7x - x$

**c**  $2x + 3x + 5y$

**d**  $8 - 6x - 2x$

**e**  $7ab + 5ba$

**f**  $3x^2 + 7x^3$

2 Remove brackets and then simplify:

**a**  $3(2x + 5) + 4(5 + 4x)$

**b**  $6 - 2(3x - 5)$

**c**  $5(2a - 3b) - 6(a - 2b)$

**d**  $3x(x^2 - 7x + 3) - (1 - 2x - 5x^2)$

3 Simplify:

a  $2x(3x)^2$

b  $\frac{3a^2b^3}{9ab^4}$

c  $\sqrt{16x^4}$

d  $(2a^2)^3 \times 3a^4$

## E LINEAR EQUATIONS AND INEQUALITIES

### EXERCISE E

**Reminder:** Multiplying or dividing both sides by a negative reverses the inequality sign.

1 Solve for  $x$ :

a  $2x + 5 = 25$

b  $3x - 7 > 11$

c  $5x + 16 = 20$

d  $\frac{x}{3} - 7 = 10$

e  $6x + 11 < 4x - 9$

f  $\frac{3x - 2}{5} = 8$

g  $1 - 2x \geq 19$

h  $\frac{1}{2}x + 1 = \frac{2}{3}x - 2$

i  $\frac{2}{3} - \frac{3x}{4} = \frac{1}{2}(2x - 1)$

2 Solve simultaneously for  $x$  and  $y$ :

a  $x + 2y = 9$   
 $x - y = 3$

b  $2x + 5y = 28$   
 $x - 2y = 2$

c  $7x + 2y = -4$   
 $3x + 4y = 14$

d  $5x - 4y = 27$   
 $3x + 2y = 9$

e  $x + 2y = 5$   
 $2x + 4y = 1$


f  $\frac{x}{2} + \frac{y}{3} = 5$   
 $\frac{x}{3} + \frac{y}{4} = 1$

## F ABSOLUTE VALUE (MODULUS)

The **modulus (absolute value)** of a real number is its size, ignoring its sign.


For example: the modulus (or absolute value) of 7 is 7, and  
the modulus (or absolute value) of  $-7$  is also 7.

Geometrically, the modulus of a real number can be interpreted as its *distance* from zero (0) on the number line. Because the modulus is distance, it cannot be negative.



Thus,

$|x|$  is the distance of  $x$  from 0 on the real number line.



**Note:**  $|x - a|$  can be considered as ‘the distance of  $x$  from  $a$ ’.

### ALGEBRAIC DEFINITION

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases} \quad \text{or} \quad |x| = \sqrt{x^2}$$

**EXERCISE F**

1 Find the value of:

**a**  $5 - (-11)$

**b**  $|5| - |-11|$

**c**  $|5 - (-11)|$

**d**  $|(-2)^2 + 11(-2)|$

**e**  $|-6| - |-8|$

**f**  $|-6 - (-8)|$

2 If  $a = -2$ ,  $b = 3$ ,  $c = -4$  find the value of:

**a**  $|a|$

**b**  $|b|$

**c**  $|a| |b|$

**d**  $|ab|$

**e**  $|a - b|$

**f**  $|a| - |b|$

**g**  $|a + b|$

**h**  $|a| + |b|$

**i**  $|a|^2$

**j**  $a^2$

**k**  $\left| \frac{c}{a} \right|$

**l**  $\frac{|c|}{|a|}$

**MODULUS EQUATIONS**It is clear that  $|x| = 2$  has two solutions,  $x = 2$  and  $x = -2$ , as  $|2| = 2$  and  $|-2| = 2$ .In general, if  $|x| = a$  where  $a > 0$ , then  $x = \pm a$ .3 Solve for  $x$ :

**a**  $|x| = 3$

**b**  $|x| = -5$

**c**  $|x| = 0$

**d**  $|x - 1| = 3$

**e**  $|3 - x| = 4$

**f**  $|x + 5| = -1$

**g**  $|3x - 2| = 1$

**h**  $|3 - 2x| = 3$

**i**  $|2 - 5x| = 12$

**G****PRODUCT EXPANSION** $y = 2(x - 1)(x + 3)$  can be expanded into the general form  $y = ax^2 + bx + c$ .Likewise,  $y = 2(x - 3)^2 + 7$  can be expanded into this form.

We will review expansion techniques.

Following is a **list of expansion rules** you should use:

- $(a + b)(c + d) = ac + ad + bc + bd$
- $(a + b)(a - b) = a^2 - b^2$  {difference of two squares}
- $(a + b)^2 = a^2 + 2ab + b^2$  }  
 $(a - b)^2 = a^2 - 2ab + b^2$  } {perfect squares}

**Example 9**

Expand and simplify:

**a**  $(2x + 1)(x + 3)$

**b**  $(3x - 2)(x + 3)$

$$\begin{aligned} \mathbf{a} \quad (2x + 1)(x + 3) \\ &= 2x^2 + 6x + x + 3 \\ &= 2x^2 + 7x + 3 \end{aligned}$$

$$\begin{aligned} \mathbf{b} \quad (3x - 2)(x + 3) \\ &= 3x^2 + 9x - 2x - 6 \\ &= 3x^2 + 7x - 6 \end{aligned}$$



**EXERCISE G**

1 Expand and simplify using  $(a + b)(c + d) = ac + ad + bc + bd$ :

- |                             |                             |                              |
|-----------------------------|-----------------------------|------------------------------|
| <b>a</b> $(2x + 3)(x + 1)$  | <b>b</b> $(3x + 4)(x + 2)$  | <b>c</b> $(5x - 2)(2x + 1)$  |
| <b>d</b> $(x + 2)(3x - 5)$  | <b>e</b> $(7 - 2x)(2 + 3x)$ | <b>f</b> $(1 - 3x)(5 + 2x)$  |
| <b>g</b> $(3x + 4)(5x - 3)$ | <b>h</b> $(1 - 3x)(2 - 5x)$ | <b>i</b> $(7 - x)(3 - 2x)$   |
| <b>j</b> $(5 - 2x)(3 - 2x)$ | <b>k</b> $-(x + 1)(x + 2)$  | <b>l</b> $-2(x - 1)(2x + 3)$ |

**Example 10**

Expand using the rule  $(a + b)(a - b) = a^2 - b^2$ :

- a**  $(5x - 2)(5x + 2)$       **b**  $(7 + 2x)(7 - 2x)$

<b>a</b> $(5x - 2)(5x + 2)$ $= (5x)^2 - 2^2$ $= 25x^2 - 4$	<b>b</b> $(7 + 2x)(7 - 2x)$ $= 7^2 - (2x)^2$ $= 49 - 4x^2$
--	--

Remember that  
 $(a + b)(a - b) = a^2 - b^2$



2 Expand using the rule  $(a + b)(a - b) = a^2 - b^2$ :

- |   |   |   |
|---|---|---|
| <b>a</b> $(x + 6)(x - 6)$               | <b>b</b> $(x + 8)(x - 8)$               | <b>c</b> $(2x - 1)(2x + 1)$               |
| <b>d</b> $(3x - 2)(3x + 2)$             | <b>e</b> $(4x + 5)(4x - 5)$             | <b>f</b> $(5x - 3)(5x + 3)$               |
| <b>g</b> $(3 - x)(3 + x)$               | <b>h</b> $(7 - x)(7 + x)$               | <b>i</b> $(7 + 2x)(7 - 2x)$               |
| <b>j</b> $(x + \sqrt{2})(x - \sqrt{2})$ | <b>k</b> $(x + \sqrt{5})(x - \sqrt{5})$ | <b>l</b> $(2x - \sqrt{3})(2x + \sqrt{3})$ |

**Example 11**

Expand using perfect square expansion rules:

- a**  $(x + 2)^2$       **b**  $(3x - 1)^2$

<b>a</b> $(x + 2)^2$ $= x^2 + 2(x)(2) + 2^2$ $= x^2 + 4x + 4$	<b>b</b> $(3x - 1)^2$ $= (3x)^2 - 2(3x)(1) + 1^2$ $= 9x^2 - 6x + 1$
---	---

Use  $(a + b)^2 = a^2 + 2ab + b^2$   
or  $(a - b)^2 = a^2 - 2ab + b^2$



3 Expand and simplify using the perfect square expansion rules:

- |                       |                       |                       |                       |
|-----------------------|-----------------------|-----------------------|-----------------------|
| <b>a</b> $(x + 5)^2$  | <b>b</b> $(x + 7)^2$  | <b>c</b> $(x - 2)^2$  | <b>d</b> $(x - 6)^2$  |
| <b>e</b> $(3 + x)^2$  | <b>f</b> $(5 + x)^2$  | <b>g</b> $(11 - x)^2$ | <b>h</b> $(10 - x)^2$ |
| <b>i</b> $(2x + 7)^2$ | <b>j</b> $(3x + 2)^2$ | <b>k</b> $(5 - 2x)^2$ | <b>l</b> $(7 - 3x)^2$ |

4 Expand the following into the general form  $y = ax^2 + bx + c$ :

- |                                 |   |  |
|---------------------------------|---|--|
| <b>a</b> $y = 2(x + 2)(x + 3)$  | <b>b</b> $y = 3(x - 1)^2 + 4$           | <b>c</b> $y = -(x + 1)(x - 7)$           |
| <b>d</b> $y = -(x + 2)^2 - 11$  | <b>e</b> $y = 4(x - 1)(x - 5)$          | <b>f</b> $y = -\frac{1}{2}(x + 4)^2 - 6$ |
| <b>g</b> $y = -5(x - 1)(x - 6)$ | <b>h</b> $y = \frac{1}{2}(x + 2)^2 - 6$ | <b>i</b> $y = -\frac{5}{2}(x - 4)^2$     |

**Example 12**

Expand and simplify:

**a**  $1 - 2(x + 3)^2$

**b**  $2(3 + x) - (2 + x)(3 - x)$

$$\begin{aligned} \mathbf{a} \quad & 1 - 2(x + 3)^2 \\ & = 1 - 2[x^2 + 6x + 9] \\ & = 1 - 2x^2 - 12x - 18 \\ & = -2x^2 - 12x - 17 \end{aligned}$$

$$\begin{aligned} \mathbf{b} \quad & 2(3 + x) - (2 + x)(3 - x) \\ & = 6 + 2x - [6 - 2x + 3x - x^2] \\ & = 6 + 2x - 6 + 2x - 3x + x^2 \\ & = x^2 + x \end{aligned}$$

The use of brackets is essential!

**5** Expand and simplify:

**a**  $1 + 2(x + 3)^2$

**c**  $3 - (3 - x)^2$

**e**  $1 + 2(4 - x)^2$

**g**  $(x + 2)^2 - (x + 1)(x - 4)$

**i**  $x^2 + 3x - 2(x - 4)^2$

**b**  $2 + 3(x - 2)(x + 3)$

**d**  $5 - (x + 5)(x - 4)$

**f**  $x^2 - 3x - (x + 2)(x - 2)$


**h**  $(2x + 3)^2 + 3(x + 1)^2$

**j**  $(3x - 2)^2 - 2(x + 1)^2$

**H****FACTORISATION**Algebraic **factorisation** is the reverse process of expansion.

For example,  $(2x + 1)(x - 3)$  is **expanded** to  $2x^2 - 5x - 3$ , whereas  $2x^2 - 5x - 3$  is **factorised** to  $(2x + 1)(x - 3)$ .

Notice that  $2x^2 - 5x - 3 = (2x + 1)(x - 3)$  has been factorised into two **linear factors**.

**Flow chart for factorising:**

**Example 13**

Fully factorise:

**a**  $3x^2 - 12x$

**b**  $4x^2 - 1$

**c**  $x^2 - 12x + 36$

**a**  $3x^2 - 12x$   
 $= 3x(x - 4)$

{has  $3x$  as common factor}

**b**  $4x^2 - 1$   
 $= (2x)^2 - 1^2$   
 $= (2x + 1)(2x - 1)$

{difference of two squares}

**c**  $x^2 - 12x + 36$   
 $= x^2 - 2(x)(6) + 6^2$   
 $= (x - 6)^2$

{perfect square form}

Remember that  
**all** factorisations  
can be checked  
by expansion!**EXERCISE H****1** Fully factorise:

**a**  $3x^2 + 9x$

**b**  $2x^2 + 7x$

**c**  $4x^2 - 10x$

**d**  $6x^2 - 15x$

**e**  $9x^2 - 25$

**f**  $16x^2 - 1$

**g**  $2x^2 - 8$

**h**  $3x^2 - 9$

**i**  $4x^2 - 20$

**j**  $x^2 - 8x + 16$

**k**  $x^2 - 10x + 25$

**l**  $2x^2 - 8x + 8$

**m**  $16x^2 + 40x + 25$

**n**  $9x^2 + 12x + 4$

**o**  $x^2 - 22x + 121$

**Example 14**

Fully factorise:

**a**  $3x^2 + 12x + 9$

**b**  $-x^2 + 3x + 10$

**a**  $3x^2 + 12x + 9$   
 $= 3(x^2 + 4x + 3)$   
 $= 3(x + 1)(x + 3)$

{has 3 as a common factor}  
{so, sum = 4, product = 3}

**b**  $-x^2 + 3x + 10$   
 $= -[x^2 - 3x - 10]$   
 $= -(x - 5)(x + 2)$

{removing  $-1$  as common factor to make  
coefficient of  $x^2$  be 1}  
{as sum =  $-3$ , product =  $-10$ }**2** Fully factorise:

**a**  $x^2 + 9x + 8$

**b**  $x^2 + 7x + 12$

**c**  $x^2 - 7x - 18$

**d**  $x^2 + 4x - 21$

**e**  $x^2 - 9x + 18$

**f**  $x^2 + x - 6$

**g**  $-x^2 + x + 2$

**h**  $3x^2 - 42x + 99$

**i**  $-2x^2 - 4x - 2$

**j**  $2x^2 + 6x - 20$

**k**  $2x^2 - 10x - 48$

**l**  $-2x^2 + 14x - 12$

**m**  $-3x^2 + 6x - 3$

**n**  $-x^2 - 2x - 1$

**o**  $-5x^2 + 10x + 40$

## FACTORISATION BY 'SPLITTING' THE $x$ -TERM

Using the distributive law to expand we see that:

$$\begin{aligned}(2x + 3)(4x + 5) &= 8x^2 + 10x + 12x + 15 \\ &= 8x^2 + 22x + 15\end{aligned}$$

We will now **reverse** the process to **factorise** the quadratic expression  $8x^2 + 22x + 15$ .

Notice that:

$$8x^2 + 22x + 15$$

*Step 1:* Split the middle term  $= 8x^2 + 10x + 12x + 15$

*Step 2:* Group in pairs  $= (8x^2 + 10x) + (12x + 15)$

*Step 3:* Factorise each pair separately  $= 2x(4x + 5) + 3(4x + 5)$

*Step 4:* Factorise fully  $= (4x + 5)(2x + 3)$

The “trick” in factorising these types of quadratic expressions is in *Step 1* where the middle term needs to be split into two so that the rest of the factorisation proceeds smoothly.

### Rules for splitting the $x$ -term:

The following procedure is recommended for factorising  $ax^2 + bx + c$ :

- find  $ac$
- find the factors of  $ac$  which add to  $b$
- if these factors are  $p$  and  $q$  replace  $bx$  by  $px + qx$
- complete the factorisation.

### Example 15

Fully factorise:

**a**  $2x^2 - x - 10$

**b**  $6x^2 - 25x + 14$

**a**  $2x^2 - x - 10$

has  $ac = 2 \times -10 = -20$ .

The factors of  $-20$  which add to  $-1$  are  $-5$  and  $+4$ .

$$\begin{aligned}&= 2x^2 - 5x + 4x - 10 \\ &= x(2x - 5) + 2(2x - 5) \\ &= (2x - 5)(x + 2)\end{aligned}$$

**b**  $6x^2 - 25x + 14$

has  $ac = 6 \times 14 = 84$ .

The factors of  $84$  which add to  $-25$  are  $-21$  and  $-4$ .

$$\begin{aligned}&= 6x^2 - 21x - 4x + 14 \\ &= 3x(2x - 7) - 2(2x - 7) \\ &= (2x - 7)(3x - 2)\end{aligned}$$

3 Fully factorise:

**a**  $2x^2 + 5x - 12$

**b**  $3x^2 - 5x - 2$

**c**  $7x^2 - 9x + 2$

**d**  $6x^2 - x - 2$

**e**  $4x^2 - 4x - 3$

**f**  $10x^2 - x - 3$

**g**  $2x^2 - 11x - 6$

**h**  $3x^2 - 5x - 28$

**i**  $8x^2 + 2x - 3$

**j**  $10x^2 - 9x - 9$

**k**  $3x^2 + 23x - 8$

**l**  $6x^2 + 7x + 2$

**m**  $-4x^2 - 2x + 6$

**n**  $12x^2 - 16x - 3$

**o**  $-6x^2 - 9x + 42$

**p**  $21x - 10 - 9x^2$

**q**  $8x^2 - 6x - 27$

**r**  $12x^2 + 13x + 3$

**s**  $12x^2 + 20x + 3$

**t**  $15x^2 - 22x + 8$

**u**  $14x^2 - 11x - 15$

### Example 16

Fully factorise:  $3(x+2) + 2(x-1)(x+2) - (x+2)^2$

$$\begin{aligned} & 3(x+2) + 2(x-1)(x+2) - (x+2)^2 \\ &= (x+2)[3 + 2(x-1) - (x+2)] \quad \{\text{as } (x+2) \text{ is the common factor}\} \\ &= (x+2)[3 + 2x - 2 - x - 2] \\ &= (x+2)(x-1) \end{aligned}$$

4 Fully factorise:

**a**  $3(x+4) + 2(x+4)(x-1)$

**b**  $8(2-x) - 3(x+1)(2-x)$

**c**  $6(x+2)^2 + 9(x+2)$

**d**  $4(x+5) + 8(x+5)^2$

**e**  $(x+2)(x+3) - (x+3)(2-x)$

**f**  $(x+3)^2 + 2(x+3) - x(x+3)$

**g**  $5(x-2) - 3(2-x)(x+7)$

**h**  $3(1-x) + 2(x+1)(x-1)$

## INVESTIGATION 1

## ANOTHER FACTORISATION TECHNIQUE



**What to do:**

1 By expanding, show that  $\frac{(ax+p)(ax+q)}{a} = ax^2 + [p+q]x + \left[\frac{pq}{a}\right]$ .

2 If  $ax^2 + bx + c = \frac{(ax+p)(ax+q)}{a}$ , show that  $p+q = b$  and  $pq = ac$ .

3 Using 2 on  $8x^2 + 22x + 15$ , we have

$$8x^2 + 22x + 15 = \frac{(8x+p)(8x+q)}{8} \quad \text{where } \begin{cases} p+q = 22 \\ pq = 8 \times 15 = 120 \end{cases}$$

So,  $p = 12$ ,  $q = 10$  (or vice versa)

$$\begin{aligned} \therefore 8x^2 + 22x + 15 &= \frac{(8x+12)(8x+10)}{8} \\ &= \frac{\cancel{4}(2x+3)\cancel{2}(4x+5)}{\cancel{8}} \\ &= (2x+3)(4x+5) \end{aligned}$$

**a** Use the method shown to factorise:

**i**  $3x^2 + 14x + 8$

**ii**  $12x^2 + 17x + 6$

**iii**  $15x^2 + 14x - 8$

**b** Check your answers to **a** using expansion.

### Example 17

Fully factorise using the ‘difference of two squares’:

**a**  $(x + 2)^2 - 9$

**b**  $(1 - x)^2 - (2x + 1)^2$

**a**  $(x + 2)^2 - 9$

$$= (x + 2)^2 - 3^2$$

$$= [(x + 2) + 3][(x + 2) - 3]$$

$$= (x + 5)(x - 1)$$

**b**  $(1 - x)^2 - (2x + 1)^2$

$$= [(1 - x) - (2x + 1)][(1 - x) + (2x + 1)]$$

$$= [1 - x - 2x - 1][1 - x + 2x + 1]$$

$$= -3x(x + 2)$$

**5** Fully factorise:

**a**  $(x + 3)^2 - 16$

**b**  $4 - (1 - x)^2$

**c**  $(x + 4)^2 - (x - 2)^2$

**d**  $16 - 4(x + 2)^2$

**e**  $(2x + 3)^2 - (x - 1)^2$

**f**  $(x + h)^2 - x^2$

**g**  $3x^2 - 3(x + 2)^2$

**h**  $5x^2 - 20(2 - x)^2$

**i**  $12x^2 - 27(3 + x)^2$

## FORMULA REARRANGEMENT

For the formula  $D = xt + p$  we say that  $D$  is the **subject**. This is because  $D$  is expressed in terms of the other variables,  $x$ ,  $t$  and  $p$ .

In formula rearrangement we require one of the other variables to be the subject.

To **rearrange** a formula we use the same processes as used for solving an equation for the variable we wish to be the subject.

### Example 18

Make  $x$  the subject of  $D = xt + p$ .

$$\text{If } D = xt + p$$

$$\text{then } xt + p = D$$

$$\therefore xt + p - p = D - p \quad \{\text{subtract } p \text{ from both sides}\}$$

$$\therefore xt = D - p$$

$$\therefore \frac{xt}{t} = \frac{D - p}{t} \quad \{\text{divide both sides by } t\}$$

$$\therefore x = \frac{D - p}{t}$$

**EXERCISE I**1 Make  $x$  the subject of:

**a**  $a + x = b$

**b**  $ax = b$

**c**  $2x + a = d$

**d**  $c + x = t$

**e**  $5x + 2y = 20$

**f**  $2x + 3y = 12$

**g**  $7x + 3y = d$

**h**  $ax + by = c$

**i**  $y = mx + c$

**Example 19**Make  $z$  the subject of  $c = \frac{m}{z}$ .

$$c = \frac{m}{z}$$

$$c \times z = \frac{m}{z} \times z \quad \{\text{multiply both sides by } z\}$$

$$\therefore cz = m$$

$$\therefore \frac{cz}{c} = \frac{m}{c} \quad \{\text{divide both sides by } c\}$$

$$\therefore z = \frac{m}{c}$$

2 Make  $z$  the subject of:

**a**  $az = \frac{b}{c}$

**b**  $\frac{a}{z} = d$

**c**  $\frac{3}{d} = \frac{2}{z}$

3 Make:

**a**  $a$  the subject of  $F = ma$

**b**  $r$  the subject of  $C = 2\pi r$

**c**  $d$  the subject of  $V = ldh$

**d**  $K$  the subject of  $A = \frac{b}{K}$

**Example 20**Make  $t$  the subject of  $s = \frac{1}{2}gt^2$  where  $t > 0$ .

$$\frac{1}{2}gt^2 = s \quad \{\text{rewrite with } t^2 \text{ on LHS}\}$$

$$\therefore 2 \times \frac{1}{2}gt^2 = 2 \times s \quad \{\text{multiply both sides by } 2\}$$

$$\therefore gt^2 = 2s$$

$$\therefore \frac{gt^2}{g} = \frac{2s}{g} \quad \{\text{divide both sides by } g\}$$

$$\therefore t^2 = \frac{2s}{g}$$

$$\therefore t = \sqrt{\frac{2s}{g}} \quad \{\text{as } t > 0\}$$

4 Make:

- a**  $r$  the subject of  $A = \pi r^2$ , ( $r > 0$ )      **b**  $x$  the subject of  $N = \frac{x^5}{a}$
- c**  $r$  the subject of  $V = \frac{4}{3}\pi r^3$       **d**  $x$  the subject of  $D = \frac{n}{x^3}$

5 Make:

- a**  $a$  the subject of  $d = \frac{\sqrt{a}}{n}$       **b**  $l$  the subject of  $T = \frac{1}{5}\sqrt{l}$
- c**  $a$  the subject of  $c = \sqrt{a^2 - b^2}$       **d**  $l$  the subject of  $T = 2\pi\sqrt{\frac{l}{g}}$
- e**  $a$  the subject of  $P = 2(a + b)$       **f**  $h$  the subject of  $A = \pi r^2 + 2\pi r h$
- g**  $r$  the subject of  $I = \frac{E}{R + r}$       **h**  $q$  the subject of  $A = \frac{B}{p - q}$

6 **a** Given the formula  $k = \frac{d^2}{2ab}$ , make  $a$  the subject of the formula.

**b** Find the value for  $a$  when  $k = 112$ ,  $d = 24$ ,  $b = 2$ .

7 The formula for determining the volume of a sphere is  $V = \frac{4}{3}\pi r^3$  where  $r$  is the radius.

**a** Make  $r$  the subject of the formula.

**b** Find the radius of a sphere having a volume of  $40 \text{ cm}^3$ .

8 The distance ( $S$  cm) travelled by an object accelerating from a stationary position is given by the formula  $S = \frac{1}{2}at^2$  where  $a$  is the acceleration ( $\text{cm/sec}^2$ ) and  $t$  is the time (seconds).

**a** Make  $t$  the subject of the formula. (Consider only  $t > 0$ .)


**b** Find the time taken for an object accelerating at  $8 \text{ cm/sec}^2$  to travel 10 m.

9 The relationship between object and image distances (in cm) for a concave mirror can

be written as  $\frac{1}{f} = \frac{1}{u} + \frac{1}{v}$  where  $f$  is the focal length,  $u$  is the object distance and  $v$  is the image distance.

**a** Make  $v$  the subject of the formula.

**b** Given a focal length of 8 cm, find the image distance for the following object distances: **i** 50 cm      **ii** 30 cm.



10 According to the theory of relativity by Einstein, the mass of a particle is given by the

formula  $m = \frac{m_0}{\sqrt{1 - \left(\frac{v}{c}\right)^2}}$ , where  $m_0$  is the mass of the particle at rest,  
 $v$  is the velocity of the particle and  
 $c$  is the velocity of light.

**a** Make  $v$  the subject of the formula, (for  $v > 0$ ).

**b** Find the velocity necessary to increase the mass of a particle to three times its rest mass, i.e.,  $m = 3m_0$ . Give the value for  $v$  as a fraction of  $c$ .

**c** A cyclotron increased the mass of an electron to  $30m_0$ . With what velocity must the electron have been travelling? [Note:  $c = 3 \times 10^8 \text{ m/s}$ ]



## J

# ADDING AND SUBTRACTING ALGEBRAIC FRACTIONS

Two or more algebraic fractions which are added (or subtracted) are combined into a single fraction by first obtaining the **least common denominator** (LCD).

For example,  $\frac{x-1}{3} - \frac{x+3}{2}$  has LCD of 6, so we write each fraction with denominator 6.

### Example 21

Write as a single fraction:     **a**  $2 + \frac{3}{x}$      **b**  $\frac{x-1}{3} - \frac{x+3}{2}$

$$\begin{aligned} \mathbf{a} \quad & 2 + \frac{3}{x} \\ &= 2 \left( \frac{x}{x} \right) + \frac{3}{x} \\ &= \frac{2x+3}{x} \end{aligned}$$

$$\begin{aligned} \mathbf{b} \quad & \frac{x-1}{3} - \frac{x+3}{2} \\ &= \frac{2}{2} \left( \frac{x-1}{3} \right) - \frac{3}{3} \left( \frac{x+3}{2} \right) \\ &= \frac{2(x-1) - 3(x+3)}{6} \\ &= \frac{2x-2-3x-9}{6} \\ &= \frac{-x-11}{6} \end{aligned}$$

### EXERCISE J

1 Write as a single fraction:

**a**  $3 + \frac{x}{5}$

**b**  $1 + \frac{3}{x}$

**c**  $3 + \frac{x-2}{2}$

**d**  $3 - \frac{x-2}{4}$

**e**  $\frac{2+x}{3} + \frac{x-4}{5}$

**f**  $\frac{2x+5}{4} - \frac{x-1}{6}$

### Example 22

Write  $\frac{3x+1}{x-2} - 2$   
as a single fraction.

$$\begin{aligned} & \frac{3x+1}{x-2} - 2 \\ &= \left( \frac{3x+1}{x-2} \right) - 2 \left( \frac{x-2}{x-2} \right) \quad \{\text{as } (x-2) \text{ is the LCD}\} \\ &= \frac{(3x+1) - 2(x-2)}{x-2} \\ &= \frac{3x+1-2x+4}{x-2} \\ &= \frac{x+5}{x-2} \end{aligned}$$

2 Write as a single fraction:

a  $1 + \frac{3}{x+2}$

b  $-2 + \frac{3}{x-4}$

c  $-3 - \frac{2}{x-1}$

d  $\frac{2x-1}{x+1} + 3$

e  $3 - \frac{x}{x+1}$

f  $-1 + \frac{4}{1-x}$

3 Write as a single fraction:

a  $\frac{3x}{2x-5} + \frac{2x+5}{x-2}$

b  $\frac{1}{x-2} - \frac{1}{x-3}$

c  $\frac{5x}{x-4} + \frac{3x-2}{x+4}$

d  $\frac{2x+1}{x-3} - \frac{x+4}{2x+1}$

# K

## CONGRUENCE AND SIMILARITY


### CONGRUENCE

Two triangles are **congruent** if they are identical in every respect apart from position, i.e., they have the same shape and size.

There are four acceptable tests for **congruence of two triangles**.



Two triangles are congruent if one of the following is true:

- corresponding sides are equal (**SSS**)
- two sides and the included angle are equal (**SAS**)
- two angles and a pair of corresponding sides are equal (**AAcorS**)
- for right angled triangles, the hypotenuses and one pair of sides are equal (**RHS**).



#### Example 23

Explain why  $\triangle ABC$  and  $\triangle DBC$  are congruent:



$\triangle$ 's ABC and DBC are congruent (SAS) as:


- $AC = DC$
- $\angle ACB = \angle DCB$ , and
- BC is common to both.

The consequence of proving congruence is that all corresponding lengths, angles and areas are equal.


**EXERCISE K.1**

- 1 Triangle ABC is isosceles with  $AC = BC$ . BC and AC are produced to E and D respectively so that  $CE = CD$ .

Prove that  $AE = BD$ .




- 2 Point P is equidistant from both AB and AC. Use congruence to show that P lies on the bisector of  $\angle BAC$ .



- 3 Two concentric circles are drawn. At P on the inner circle a tangent is drawn and it meets the other circle at A and B.

Use triangle congruence to prove that P is the midpoint of AB.





**SIMILARITY**

Two triangles are **similar** if one is an enlargement of the other. Consequently, similar triangles are **equiangular**. Similar triangles have corresponding sides in the same **ratio**.

**Example 24**

Establish that a pair of triangles is similar and find  $x$  if  $BD = 20$  cm:









$\alpha$	$\beta$	$\bullet$	
-	$x + 2$	$x$	small $\Delta$
-	20	12	large $\Delta$

The triangles are equiangular and hence similar.

$$\begin{aligned} \therefore \frac{x + 2}{20} &= \frac{x}{12} && \{\text{same ratio}\} \\ \therefore 12(x + 2) &= 20x \\ \therefore 12x + 24 &= 20x \\ \therefore 24 &= 8x \\ \therefore x &= 3 \end{aligned}$$

**EXERCISE K.2**

1 In the following, establish that a pair of triangles is similar, and find  $x$ :



2 A father and son are standing side-by-side. How tall is the son if the father is 1.8 m tall and casts a shadow 3.2 m long, while his son's shadow is 2.4 m long?

**L****COORDINATE GEOMETRY****THE NUMBER PLANE**

The position or location of any point in the **number plane** can be specified in terms of an **ordered pair** of numbers  $(x, y)$ , where:

$x$  is the **horizontal step** from a fixed point  $O$ , and  $y$  is the **vertical step** from  $O$ .

Once an **origin**  $O$ , has been given, two perpendicular axes are drawn.

The  $x$ -axis is horizontal and the  $y$ -axis is vertical.


The **number plane** is also known as either:

- the **2-dimensional plane**, or
- the **Cartesian plane**, (named after **René Descartes**).

**Note:**  $(a, b)$  is called an **ordered pair**, where  $a$  and  $b$  are often referred to as **the coordinates** of the point.

$a$  is called the  $x$ -coordinate and

$b$  is called the  $y$ -coordinate.

**THE DISTANCE FORMULA**

If  $A(x_1, y_1)$  and  $B(x_2, y_2)$  are two points in a plane, then the distance  $d$ ,

between these points is given by 
$$AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

**Example 25**


Find the distance between  $A(-2, 1)$  and  $B(3, 4)$ .

$$\begin{array}{ccc}
 A(-2, 1) & B(3, 4) & \\
 \uparrow & \uparrow & \\
 x_1 & y_1 & x_2 & y_2
 \end{array}
 \quad
 \begin{aligned}
 AB &= \sqrt{(3 - (-2))^2 + (4 - 1)^2} \\
 &= \sqrt{5^2 + 3^2} \\
 &= \sqrt{25 + 9} \\
 &= \sqrt{34} \text{ units}
 \end{aligned}$$

This distance formula saves us having to graph the points each time we want to find a distance.

**THE MIDPOINT FORMULA**

If  $M$  is halfway between points  $A$  and  $B$  then  $M$  is the **midpoint** of  $AB$ .



If  $A(x_1, y_1)$  and  $B(x_2, y_2)$  are two points then the **midpoint**  $M$  of  $AB$  has coordinates  $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$ .

**Example 26**

Find the coordinates of the midpoint of  $AB$  for  $A(-1, 3)$  and  $B(4, 7)$ .

$$x\text{-coordinate of midpoint} = \frac{-1 + 4}{2} = \frac{3}{2} = 1\frac{1}{2}$$

$$y\text{-coordinate of midpoint} = \frac{3 + 7}{2} = 5$$

$\therefore$  the midpoint of  $AB$  is  $(1\frac{1}{2}, 5)$

**SLOPE (OR GRADIENT) OF A LINE**

When looking at line segments drawn on a set of axes, it is clear that different line segments are inclined to the horizontal at different angles, i.e., some appear to be steeper than others.

The **slope** or **gradient** of a line is a measure of its steepness.

If  $A$  is  $(x_1, y_1)$  and  $B$  is  $(x_2, y_2)$  then the **slope** of  $AB$  is  $\frac{y_2 - y_1}{x_2 - x_1}$ .


**Example 27**


Find the slope of the line through  $(3, -2)$  and  $(6, 4)$ .

$$\begin{array}{ccc}
 (3, -2) & (6, 4) & \\
 \uparrow & \uparrow & \\
 x_1 & y_1 & x_2 & y_2
 \end{array}
 \quad
 \text{slope} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - (-2)}{6 - 3} = 2$$

**Note:** • **horizontal lines** have a gradient of **0 (zero)**

• **vertical lines** have an **undefined** gradient.

•  forward sloping lines have **positive** gradients

•  backward sloping lines have **negative** gradient

• **parallel lines** have equal gradients

• the slopes of **perpendicular lines** are *negative reciprocals*

i.e., if the slopes are  $m_1$  and  $m_2$  then  $m_2 = \frac{-1}{m_1}$  or  $m_1 m_2 = -1$ .

**Note:**  $m_1 m_2 = -1$  is not true if the lines are parallel to the axes.

## EQUATIONS OF LINES

The **equation of a line** is an equation which states the connection between the  $x$  and  $y$  values for every point on the line, and only for points on the line.

Equations of lines have various forms:

- All **vertical lines** have equations of the form  $x = a$ , ( $a$  is a constant).
- All **horizontal lines** have equations of the form  $y = c$ , ( $c$  is a constant).
- If a straight line has gradient (or slope)  $m$  and passes through  $(a, b)$  then it has equation

$$\frac{y - b}{x - a} = m \quad \text{or} \quad y - b = m(x - a) \quad \{\text{point-gradient form}\}$$

which can be rearranged into  $y = mx + c$  **{slope-intercept form}**  
or  $Ax + By = C$  **{general form}**

### Example 28

Find, in *slope-intercept form*, the equation of the line through  $(-1, 3)$  with a slope of 5.

The equation of the line is  $\frac{y - 3}{x - (-1)} = 5$  i.e.,  $\frac{y - 3}{x + 1} = 5$

$$\therefore y - 3 = 5(x + 1)$$

$$\therefore y - 3 = 5x + 5$$

$$\therefore y = 5x + 8$$

To find the equation of a line we need to know its gradient (or slope) and a point on it.



### Example 29

Find, in *general form*, the equation of the line through  $(1, -5)$  and  $(5, -2)$ .

$$\begin{aligned} \text{The slope} &= \frac{-2 - (-5)}{5 - 1} \\ &= \frac{3}{4} \end{aligned}$$

So, the equation is

$$\frac{y - (-2)}{x - 5} = \frac{3}{4}$$

$$\text{i.e., } \frac{y + 2}{x - 5} = \frac{3}{4}$$


$$\therefore 4y + 8 = 3x - 15$$

$$\therefore 3x - 4y = 23$$

## INTERCEPTS

Axis **intercepts** are the  $x$ - and  $y$ -values where a graph cuts the coordinate axes.

The  $x$ -intercept is found by letting  $y = 0$ .  
The  $y$ -intercept is found by letting  $x = 0$ .



### Example 30

For the line with equation  $2x - 3y = 12$ , find axis intercepts.

For  $2x - 3y = 12$ ,

$$\begin{array}{ll} \text{when } x = 0, & -3y = 12 \\ & \therefore y = -4 \end{array} \qquad \begin{array}{ll} \text{when } y = 0, & 2x = 12 \\ & \therefore x = 6 \end{array}$$

## DOES A POINT LIE ON A LINE?

A point lies on a line if its coordinates satisfy the equation of the line.

### Example 31

Does  $(3, -2)$  lie on the line with equation  $5x - 2y = 20$ ?

Substituting  $(3, -2)$  into  $5x - 2y = 20$  gives  
 $5(3) - 2(-2) = 20$   
 i.e.,  $19 = 20$  which is false

$\therefore (3, -2)$  does not lie on the line.

## WHERE GRAPHS MEET

### Example 32

Use graphical methods to find where the lines  $x + y = 6$  and  $2x - y = 6$  meet.

For  $x + y = 6$

when  $x = 0$ ,  $y = 6$

when  $y = 0$ ,  $x = 6$

$x$	0	6
$y$	6	0

For  $2x - y = 6$

when  $x = 0$ ,  $-y = 6$ ,

$\therefore y = -6$


when  $y = 0$ ,  $2x = 6$ ,

$\therefore x = 3$


$x$	0	3
$y$	-6	0

The graphs meet at  $(4, 2)$ .

Check:  $4 + 2 = 6$  ✓ and  $2 \times 4 - 2 = 6$  ✓



There are three possible situations which may occur. These are:



The lines meet in a single **point of intersection**.



The lines are **parallel** and **never meet**. So, there is no point of intersection.



The lines are **coincident** (the same line) and so there are infinitely many points of intersection.

## INVESTIGATION 2 FINDING WHERE LINES MEET USING TECHNOLOGY



**Graphing packages** and **graphics calculators** can be used to plot straight line graphs and hence find the point of intersection of the straight lines. This can be useful if the solutions are not integer values, although an algebraic method can also be used. However, most graphing packages and graphics calculators require the equation to be entered in the form  $y = mx + c$ .

Consequently, if an equation is given in **general form**, it must be rearranged into **slope-intercept form**. For example, if we wish to use technology to find the point of intersection of  $4x + 3y = 10$  and  $x - 2y = -3$ :

*Step 1:* We **rearrange** each equation into the form  $y = mx + c$ , i.e.,


$$\begin{array}{ll} 4x + 3y = 10 & \text{and } x - 2y = -3 \\ \therefore 3y = -4x + 10 & \therefore -2y = -x - 3 \\ \therefore y = -\frac{4}{3}x + \frac{10}{3} & \therefore y = \frac{x}{2} + \frac{3}{2} \end{array}$$

*Step 2:* If you are using the **graphing package**, click on the icon to open the package and enter the two equations.

If you are using a **graphics calculator**, enter the functions  $Y_1 = -4X/3 + 10/3$  and  $Y_2 = X/2 + 3/2$ .



*Step 3:* Draw the **graphs** of the functions on the same set of axes. (You may have to change the viewing **window** if using a graphics calculator.)



*Step 4:* Use the built in functions to calculate the point of **intersection**. Thus, the point of intersection is (1, 2).

### What to do:

**1** Use technology to find the point of intersection of:

**a**  $y = x + 4$   
 $5x - 3y = 0$

**b**  $x + 2y = 8$   
 $y = 7 - 2x$

**c**  $x - y = 5$   
 $2x + 3y = 4$

**d**  $2x + y = 7$   
 $3x - 2y = 1$

**e**  $y = 3x - 1$   
 $3x - y = 6$

**f**  $y = -\frac{2x}{3} + 2$   
 $2x + 3y = 6$


**2** Comment on the use of technology to find the point(s) of intersection in **1 e** and **1 f**.

**Note:** Whilst you have not yet studied many functions, you should be able to graph given functions (usually curves) and find where they meet.




**EXERCISE L**

- 1 Use the distance formula to find the distance between the following pairs of points:
- a** A(1, 3) and B(4, 5)                      **b** O(0, 0) and C(3, -5)  
**c** P(5, 2) and Q(1, 4)                      **d** S(0, -3) and T(-1, 0)
- 2 Find the midpoint of AB for:
- a** A(3, 6) and B(1, 0)                      **b** A(5, 2) and B(-1, -4)  
**c** A(7, 0) and B(0, 3)                      **d** A(5, -2) and B(-1, -3)
- 3 By finding an appropriate  $y$ -step and  $x$ -step, determine the slope of each of the following lines:



- 4 Find the slope of the line passing through:
- a** (2, 3) and (4, 7)                      **b** (3, 2) and (5, 8)  
**c** (-1, 2) and (-1, 5)                      **d** (4, -3) and (-1, -3)  
**e** (0, 0) and (-1, 4)                      **f** (3, -1) and (-1, -2)
- 5 Classify the following pairs of lines as parallel, perpendicular or neither. Give a reason.



- 6 State the slope of the line which is perpendicular to the line with slope:
- a**  $\frac{3}{4}$                       **b**  $\frac{11}{3}$                       **c** 4                      **d**  $-\frac{1}{3}$                       **e** -5                      **f** 0
- 7 Find the equation of the line in slope-intercept form, through:
- a** (4, 1) with slope 2                      **b** (1, 2) with slope -2                      **c** (5, 0) with slope 3  
**d** (-1, 7) with slope -3                      **e** (1, 5) with slope -4                      **f** (2, 7) with slope 1

8 Find the equation of the line, in general form, through:

- a** (2, 1) with slope  $\frac{3}{2}$     **b** (1, 4) with slope  $-\frac{3}{2}$     **c** (4, 0) with slope  $\frac{1}{3}$   
**d** (0, 6) with slope  $-4$     **e**  $(-1, -3)$  with slope 3    **f** (4,  $-2$ ) with slope  $-\frac{4}{9}$

9 Find the equations of the lines through:

- a** (0, 1) and (3, 2)                      **b** (1, 4) and (0,  $-1$ )  
**c** (2,  $-1$ ) and  $(-1, -4)$                 **d** (0,  $-2$ ) and (5, 2)  
**e** (3, 2) and  $(-1, 0)$                     **f**  $(-1, -1)$  and (2,  $-3$ )

10 Find the equations of the lines through:

- a** (3,  $-2$ ) and (5,  $-2$ )    **b** (6, 7) and (6,  $-11$ )    **c**  $(-3, 1)$  and  $(-3, -3)$

11 Copy and complete:

	Equation of line	Slope	$x$ -intercept	$y$ -intercept
<b>a</b>	$2x - 3y = 6$			
<b>b</b>	$4x + 5y = 20$			
<b>c</b>	$y = -2x + 5$			
<b>d</b>	$x = 8$			
<b>e</b>	$y = 5$			
<b>f</b>	$x + y = 11$			
<b>g</b>	$4x + y = 8$			
<b>h</b>	$x - 3y = 12$			

If a line has equation  $y = mx + c$  then the slope of the line is 'm'.



12 **a** Does (3, 4) lie on the line with equation  $3x - 2y = 1$ ?

**b** Does  $(-2, 5)$  lie on the line with equation  $5x + 3y = -5$ ?

**c** Does  $(6, -\frac{1}{2})$  lie on the line  $3x - 8y = 22$ ?

13 Use graphical methods to find where the following lines meet:

**a**  $x + 2y = 8$   
 $y = 2x - 6$

**b**  $y = -3x - 3$   
 $3x - 2y = -12$

**c**  $3x + y = -3$   
 $2x - 3y = -24$

**d**  $2x - 3y = 8$   
 $3x + 2y = 12$

**e**  $x + 3y = 10$   
 $2x + 6y = 11$

**f**  $5x + 3y = 10$   
 $10x + 6y = 20$

### Example 33


A straight road is to pass through points A(5, 3) and B(1, 8).

**a** Find where this road meets the road given by:

- i**  $x = 3$     **ii**  $y = 4$

**b** If we wish to refer to points on road AB, but between A and B, how can we indicate this?

**c** Does C(23,  $-20$ ) lie on the road?



First we must find the equation of the line representing the road.

Its slope is  $m = \frac{3 - 8}{5 - 1} = -\frac{5}{4}$

So, its equation is  $\frac{y - 3}{x - 5} = -\frac{5}{4}$

i.e.,  $4(y - 3) = -5(x - 5)$

i.e.,  $4y - 12 = -5x + 25$

i.e.,  $5x + 4y = 37$

**a i** when  $x = 3$ ,  $5(3) + 4y = 37$

$\therefore 15 + 4y = 37$

$\therefore 4y = 22$

$\therefore y = 5\frac{1}{2}$

i.e., meets at  $(3, 5\frac{1}{2})$

**ii** when  $y = 4$ ,  $5x + 4(4) = 37$

$\therefore 5x + 16 = 37$

$\therefore 5x = 21$

$\therefore x = 4.2$

$\therefore$  meets at  $(4.2, 4)$

**b** We state the possible  $x$ -value restriction i.e.,  $1 \leq x \leq 5$ .

**c** If  $C(23, -20)$  lies on the line, its coordinates must satisfy the line's equation.

Now LHS =  $5(23) + 4(-20)$

=  $115 - 80$

=  $35$

$\neq 37 \therefore C$  does not lie on the road.

**14** Find the equation of the:

**a** horizontal line through  $(3, -4)$

**c** vertical line through  $(-1, -3)$

**e**  $x$ -axis

**b** vertical line with  $x$ -intercept 5

**d** horizontal line with  $y$ -intercept 2

**f**  $y$ -axis

**15** Find the equation of the line which is:

**a** through  $A(-1, 4)$  and with slope  $\frac{3}{4}$

**b** through  $P(2, -5)$  and  $Q(7, 0)$

**c** parallel to the line with equation  $y = 3x - 2$ , but passes through  $(0, 0)$

**d** parallel to the line with equation  $2x + 3y = 8$ , but passes through  $(-1, 7)$

**e** perpendicular to the line with equation  $y = -2x + 5$  and passes through  $(3, -1)$

**f** perpendicular to the line with equation  $3x - y = 11$  and passes through  $(-2, 5)$ .


**16** A is the town hall on Scott Street and D is a Post Office on Keach Avenue. Diagonal Road intersects Scott Street at B and Keach Avenue at C.

**a** Find the equation of Keach Avenue.

**b** Find the equation of Peacock Street.


**c** Find the equation of Diagonal Road. (Be careful!)

**d** Plunkit Street lies on map reference line  $x = 8$ . Where does Plunkit Street intersect Keach Avenue?



**Example 34**

Find the equation of the tangent to the circle with centre (2, 3) at the point (-1, 5).



$$\text{slope of CP is } \frac{3 - 5}{2 - (-1)} = \frac{-2}{3} = -\frac{2}{3}$$

$\therefore$  the slope of the tangent at P is  $\frac{3}{2}$ ,  
and since the tangent is through (-1, 5)  
the equation is

$$\frac{y - 5}{x - (-1)} = \frac{3}{2}$$

$$\therefore 2(y - 5) = 3(x + 1)$$

$$\therefore 2y - 10 = 3x + 3$$

$$\text{i.e., } 3x - 2y = -13$$



The tangent is perpendicular to the radius at the point of contact.

**17** Find the equation of the tangent to the circle with centre:


**a** (0, 2) at (-1, 5)

**b** (3, -1) at (-1, 1)

**c** (2, -2) at (5, -2)

**Example 35**

Mining towns are situated at B(1, 6) and A(5, 2). Where should a railway siding S be located so that ore trucks from either A or B would travel equal distances to a railway line with equation  $x = 11$ ?



We let a general point on the line  $x = 11$  have coordinates (11,  $a$ ) say.

Now  $BS = AS$

$$\therefore \sqrt{(11 - 1)^2 + (a - 6)^2} = \sqrt{(11 - 5)^2 + (a - 2)^2}$$

$$\therefore 10^2 + (a - 6)^2 = 6^2 + (a - 2)^2 \quad \{\text{squaring both sides}\}$$

$$\therefore 100 + a^2 - 12a + 36 = 36 + a^2 - 4a + 4$$


$$\therefore -12a + 4a = 4 - 100$$

$$\therefore -8a = -96$$

$$\therefore a = 12$$


So, the railway siding should be located at (11, 12).

**18** A(5, 5) and B(7, 10) are houses and  $y = 8$  is a gas pipeline. Where should the one outlet from the pipeline be placed so that it is the same distance from both houses so they pay equal service costs?




**19** CD is a water pipeline. A and B are two towns. A pumping station is to be located on the pipeline to pump water to A and B. Each town is to pay for their own service pipes and insist on equality of costs.

- a Where should C be located for equality of costs to occur?
- b What is the total length of service pipe required?
- c If the towns agree to pay equal amounts, would it be cheaper to install the service pipeline from D to B to A?



**Example 36**



A tunnel through the mountains connects town Q(2, 4) to the port at P.

P is on grid reference  $x = 6$  and the distance between the town and the port is 5 km.

Assuming the diagram is reasonably accurate, what is the horizontal grid reference of the port?

Let P be (6, a) say.

$$\begin{aligned}
 \text{Now } PQ &= 5 \\
 \therefore \sqrt{(6-2)^2 + (a-4)^2} &= 5 \\
 \therefore \sqrt{16 + (a-4)^2} &= 5 \\
 \therefore 16 + (a-4)^2 &= 25 \\
 \therefore (a-4)^2 &= 9 \\
 \therefore a-4 &= \pm 3 \\
 \therefore a &= 4 \pm 3 = 7 \text{ or } 1
 \end{aligned}$$

But P is further North than Q  $\therefore a > 4$

So P is at (6, 7) and the horizontal grid reference is  $y = 7$ .

**20**  $y = 8$  Clifton Highway

Jason's girlfriend lives in a house on Clifton Highway which has equation  $y = 8$ . The distance 'as the crow flies' from Jason's house to his girlfriend's house is 11.73 km. If Jason lives at (4, 1), what are the coordinates of his girlfriend's house?



Scale: 1 unit  $\equiv$  1 km.

- 21** A circle has centre  $(a, b)$  and radius  $r$  units.  $P(x, y)$  moves on the circle.

Show that  $(x - a)^2 + (y - b)^2 = r^2$ .


- 22**  $(x - a)^2 + (y - b)^2 = r^2$  is the equation of a circle centre  $(a, b)$ , radius  $r$ .

Find the equation of the circle with:

- a** centre  $(4, 3)$  and radius 5 units
  - b** centre  $(-1, 5)$  and radius 2 units
  - c** centre  $(0, 0)$  and radius 10 units
  - d** ends of a diameter  $(-1, 5)$  and  $(3, 1)$
- 23** Find the centre and radius of the circle:
- a**  $(x-1)^2 + (y-3)^2 = 4$
  - b**  $x^2 + (y+2)^2 = 16$
  - c**  $x^2 + y^2 = 7$
- 24** Consider the circle with equation  $(x - 2)^2 + (y + 3)^2 = 20$ .
- a** State the circle's centre and radius.
  - b** Show that  $(4, 1)$  lies on the circle.
  - c** Find the equation of the tangent to the circle at  $(4, 1)$ .

- 25** The perpendicular bisector of a chord of a circle passes through its centre.

Find the centre of a circle passing through points  $P(5, 7)$ ,  $Q(7, 1)$  and  $R(-1, 5)$  by finding the perpendicular bisectors of  $PQ$  and  $QR$  and solving them simultaneously.



**EXERCISE A**


- 1 a**  $\sqrt{15}$  **b** 3 **c** 4 **d** 12 **e** 42 **f**  $\sqrt{6}$  **g**  $\sqrt{2}$  **h**  $\sqrt{6}$   
**2 a**  $5\sqrt{2}$  **b**  $-\sqrt{2}$  **c**  $2\sqrt{5}$  **d**  $8\sqrt{5}$  **e**  $-2\sqrt{5}$   
**f**  $9\sqrt{3}$  **g**  $-3\sqrt{6}$  **h**  $3\sqrt{2}$   
**3 a**  $2\sqrt{2}$  **b**  $2\sqrt{3}$  **c**  $2\sqrt{5}$  **d**  $4\sqrt{2}$  **e**  $3\sqrt{3}$  **f**  $3\sqrt{5}$   
**g**  $4\sqrt{3}$  **h**  $3\sqrt{6}$  **i**  $5\sqrt{2}$  **j**  $4\sqrt{5}$  **k**  $4\sqrt{6}$  **l**  $6\sqrt{3}$   
**4 a**  $2\sqrt{3}$  **b**  $8\sqrt{2}$  **c**  $5\sqrt{6}$  **d**  $10\sqrt{3}$  **e**  $3\sqrt{3}$  **f**  $-\sqrt{2}$   
**5 a**  $\frac{\sqrt{2}}{2}$  **b**  $2\sqrt{3}$  **c**  $\frac{7\sqrt{2}}{2}$  **d**  $2\sqrt{5}$  **e**  $5\sqrt{2}$  **f**  $3\sqrt{6}$   
**g**  $4\sqrt{3}$  **h**  $\frac{5\sqrt{7}}{7}$  **i**  $2\sqrt{7}$  **j**  $\sqrt{6}$

**EXERCISE B**

- 1 a**  $2.59 \times 10^2$  **b**  $2.59 \times 10^5$  **c**  $2.59 \times 10^0$   
**d**  $2.59 \times 10^{-1}$  **e**  $2.59 \times 10^{-4}$  **f**  $4.07 \times 10^1$   
**g**  $4.07 \times 10^3$  **h**  $4.07 \times 10^{-2}$  **i**  $4.07 \times 10^5$   
**j**  $4.07 \times 10^8$  **k**  $4.07 \times 10^{-5}$   
**2 a**  $1.495 \times 10^{11}$  m **b**  $3 \times 10^{-4}$  mm **c**  $1 \times 10^{-3}$  mm  
**d**  $1.5 \times 10^7$  °C **e**  $3 \times 10^5$   
**3 a** 4000 **b** 500 **c** 2100 **d** 78 000 **e** 380 000  
**f** 86 **g** 43 300 000 **h** 60 000 000  
**4 a** 0.004 **b** 0.05 **c** 0.0021 **d** 0.000 78  
**e** 0.000 038 **f** 0.86 **g** 0.000 000 433 **h** 0.000 000 6  
**5 a** 0.000 000 9 m **b** 6 130 000 000 **c** 100 000 light years  
**d** 0.000 01 mm  
**6 a**  $1.64 \times 10^{10}$  **b**  $4.12 \times 10^{-3}$  **c**  $5.27 \times 10^{-18}$   
**d**  $1.36 \times 10^2$  **e**  $2.63 \times 10^{-6}$  **f**  $1.73 \times 10^9$   
**7 a**  $1.30 \times 10^5$  km **b**  $9.07 \times 10^5$  km **c**  $9.47 \times 10^7$  km  
**8 a**  $1.8 \times 10^{10}$  m **b**  $2.59 \times 10^{13}$  m **c**  $9.47 \times 10^{15}$  m

**EXERCISE C**

- 1 a** The set of all  $x$  such that  $x$  is greater than 5.  
**b** The set of all  $x$  such that  $x$  is less than or equal to 3.  
**c** The set of all  $y$  such that  $y$  lies between 0 and 6.  
**d** The set of all  $x$  such that  $x$  is greater than or equal to 2, but less than or equal to 4.  
**e** The set of all  $t$  such that  $t$  lies between 1 and 5.  
**f** The set of all  $n$  such that  $n$  is less than 2 or greater than or equal to 6.  
**2 a**  $\{x : x > 2\}$  **b**  $\{x : 1 < x \leq 5\}$   
**c**  $\{x : x \leq -2 \text{ or } x \geq 3\}$  **d**  $\{x : x \in \mathbb{Z}, -1 \leq x \leq 3\}$   
**e**  $\{x : x \in \mathbb{Z}, 0 \leq x \leq 5\}$  **f**  $\{x : x < 0\}$


**EXERCISE D**

- 1 a**  $10x - 10$  **b**  $9x$  **c**  $5x + 5y$  **d**  $8 - 8x$  **e**  $12ab$   
**f** cannot be simplified  
**2 a**  $22x + 35$  **b**  $16 - 6x$  **c**  $4a - 3b$

- d**  $3x^3 - 16x^2 + 11x - 1$   
**3 a**  $18x^3$  **b**  $\frac{a}{3b}$  **c**  $4x^2$  **d**  $24a^{10}$

**EXERCISE E**

- 1 a**  $x = 10$  **b**  $x > 6$  **c**  $x = \frac{4}{5}$  **d**  $x = 51$  **e**  $x < -10$   
**f**  $x = 14$  **g**  $x \leq -9$  **h**  $x = 18$  **i**  $x = \frac{2}{3}$   
**2 a**  $x = 5, y = 2$  **b**  $x = \frac{22}{3}, y = \frac{8}{3}$  **c**  $x = -2, y = 5$   
**d**  $x = \frac{45}{11}, y = -\frac{18}{11}$  **e** no solution  
**f**  $x = 66, y = -84$

**EXERCISE F**

- 1 a** 16 **b** -6 **c** 16 **d** 18 **e** -2 **f** 2  
**2 a** 2 **b** 3 **c** 6 **d** 6 **e** 5 **f** -1 **g** 1 **h** 5  
**i** 4 **j** 4 **k** 2 **l** 2  
**3 a**  $x = \pm 3$  **b** no solution **c**  $x = 0$  **d**  $x = 4$  or  $-2$   
**e**  $x = -1$  or  $7$  **f** no solution **g**  $x = 1$  or  $\frac{1}{3}$   
**h**  $x = 0$  or  $3$  **i**  $x = -2$  or  $\frac{14}{5}$

**EXERCISE G**

- 1 a**  $2x^2 + 5x + 3$  **b**  $3x^2 + 10x + 8$  **c**  $10x^2 + x - 2$   
**d**  $3x^2 + x - 10$  **e**  $-6x^2 + 17x + 14$  **f**  $-6x^2 - 13x + 5$   
**g**  $15x^2 + 11x - 12$  **h**  $15x^2 - 11x + 2$  **i**  $2x^2 - 17x + 21$   
**j**  $4x^2 - 16x + 15$  **k**  $-x^2 - 3x - 2$  **l**  $-4x^2 - 2x + 6$   
**2 a**  $x^2 - 36$  **b**  $x^2 - 64$  **c**  $4x^2 - 1$  **d**  $9x^2 - 4$   
**e**  $16x^2 - 25$  **f**  $25x^2 - 9$  **g**  $9 - x^2$  **h**  $49 - x^2$   
**i**  $49 - 4x^2$  **j**  $x^2 - 2$  **k**  $x^2 - 5$  **l**  $4x^2 - 3$   
**3 a**  $x^2 + 10x + 25$  **b**  $x^2 + 14x + 49$  **c**  $x^2 - 4x + 4$   
**d**  $x^2 - 12x + 36$  **e**  $x^2 + 6x + 9$  **f**  $x^2 + 10x + 25$   
**g**  $x^2 - 22x + 121$  **h**  $x^2 - 20x + 100$  **i**  $4x^2 + 28x + 49$   
**j**  $9x^2 + 12x + 4$  **k**  $4x^2 - 20x + 25$  **l**  $9x^2 - 42x + 49$   
**4 a**  $y = 2x^2 + 10x + 12$  **b**  $y = 3x^2 - 6x + 7$   
**c**  $y = -x^2 + 6x + 7$  **d**  $y = -x^2 - 4x - 15$   
**e**  $y = 4x^2 - 24x + 20$  **f**  $y = -\frac{1}{2}x^2 - 4x - 14$   
**g**  $y = -5x^2 + 35x - 30$  **h**  $y = \frac{1}{2}x^2 + 2x - 4$   
**i**  $y = -\frac{5}{2}x^2 + 20x - 40$   
**5 a**  $2x^2 + 12x + 19$  **b**  $3x^2 + 3x - 16$  **c**  $-x^2 + 6x - 6$   
**d**  $-x^2 - x + 25$  **e**  $2x^2 - 16x + 33$  **f**  $-3x + 4$   
**g**  $7x + 8$  **h**  $7x^2 + 18x + 12$  **i**  $-x^2 + 19x - 32$   
**j**  $7x^2 - 16x + 2$

**EXERCISE H**

- 1 a**  $3x(x+3)$  **b**  $x(2x+7)$  **c**  $2x(2x-5)$  **d**  $3x(2x-5)$   
**e**  $(3x-5)(3x+5)$  **f**  $(4x+1)(4x-1)$  **g**  $2(x-2)(x+2)$   
**h**  $3(x+\sqrt{3})(x-\sqrt{3})$  **i**  $4(x+\sqrt{5})(x-\sqrt{5})$  **j**  $(x-4)^2$   
**k**  $(x-5)^2$  **l**  $2(x-2)^2$  **m**  $(4x+5)^2$  **n**  $(3x+2)^2$   
**o**  $(x-11)^2$   
**2 a**  $(x+8)(x+1)$  **b**  $(x+4)(x+3)$  **c**  $(x-9)(x+2)$   
**d**  $(x+7)(x-3)$  **e**  $(x-6)(x-3)$  **f**  $(x+3)(x-2)$   
**g**  $-(x-2)(x+1)$  **h**  $3(x-11)(x-3)$  **i**  $-2(x+1)^2$   
**j**  $2(x+5)(x-2)$  **k**  $2(x-8)(x+3)$  **l**  $-2(x-6)(x-1)$   
**m**  $-3(x-1)^2$  **n**  $-(x+1)^2$  **o**  $-5(x-4)(x+2)$   
**3 a**  $(2x-3)(x+4)$  **b**  $(3x+1)(x-2)$  **c**  $(7x-2)(x-1)$   
**d**  $(3x-2)(2x+1)$  **e**  $(2x-3)(2x+1)$  **f**  $(5x-3)(2x+1)$   
**g**  $(2x+1)(x-6)$  **h**  $(3x+7)(x-4)$  **i**  $(4x+3)(2x-1)$   
**j**  $(5x+3)(2x-3)$  **k**  $(3x-1)(x+8)$  **l**  $(3x+2)(2x+1)$   
**m**  $-2(2x+3)(x-1)$  **n**  $(6x+1)(2x-3)$   
**o**  $-3(2x+7)(x-2)$  **p**  $-(3x-2)(3x-5)$

**q**  $(4x-9)(2x+3)$  **r**  $(4x+3)(3x+1)$   
**s**  $(6x+1)(2x+3)$  **t**  $(5x-4)(3x-2)$   
**u**  $(7x+5)(2x-3)$

**4 a**  $(x+4)(2x+1)$  **b**  $(2-x)(5-3x)$  **c**  $3(x+2)(2x+7)$   
**d**  $4(x+5)(2x+11)$  **e**  $2x(x+3)$  **f**  $5(x+3)$   
**g**  $(x-2)(3x+26)$  **h**  $(x-1)(2x-1)$

**5 a**  $(x+7)(x-1)$  **b**  $(x+1)(3-x)$  **c**  $12(x+1)$   
**d**  $-4x(x+4)$  **e**  $(3x+2)(x+4)$  **f**  $h(2x+h)$   
**g**  $-12(x+1)$  **h**  $-5(3x-4)(x-4)$  **i**  $-3(x+9)(5x+9)$

**EXERCISE I**

**1 a**  $x = b - a$  **b**  $x = \frac{b}{a}$  **c**  $x = \frac{d-a}{2}$  **d**  $x = t - c$

**e**  $x = \frac{20-2y}{5}$  **f**  $x = \frac{12-3y}{2}$  **g**  $x = \frac{d-3y}{7}$

**h**  $x = \frac{c-by}{a}$  **i**  $x = \frac{y-c}{m}$

**2 a**  $z = \frac{b}{ac}$  **b**  $z = \frac{a}{d}$  **c**  $z = \frac{2d}{3}$

**3 a**  $a = \frac{F}{m}$  **b**  $r = \frac{C}{2\pi}$  **c**  $d = \frac{V}{lh}$  **d**  $K = \frac{b}{A}$

**4 a**  $r = \sqrt{\frac{A}{\pi}}$  **b**  $x = \sqrt[5]{aN}$  **c**  $r = \sqrt[3]{\frac{3V}{4\pi}}$  **d**  $x = \sqrt[3]{\frac{n}{D}}$

**5 a**  $a = d^2n^2$  **b**  $l = 25T^2$  **c**  $a = \pm\sqrt{b^2+c^2}$  **d**  $l = \frac{gT^2}{4\pi^2}$

**e**  $a = \frac{P}{2} - b$  **f**  $h = \frac{A - \pi r^2}{2\pi r}$  **g**  $r = \frac{E}{I} - R$  **h**  $q = p - \frac{B}{A}$

**6 a**  $a = \frac{d^2}{2kb}$  **b** 1.29 **7 a**  $r = \sqrt[3]{\frac{3V}{4\pi}}$  **b** 2.122 cm

**8 a**  $t = \sqrt{\frac{2S}{a}}$  **b** 15.81 sec

**9 a**  $v = \frac{uf}{u-f}$  **b i** 9.52 cm **ii** 10.9 cm

**10 a**  $v = \sqrt{c^2 \left(1 - \frac{m_0^2}{m^2}\right)} = \frac{c}{m} \sqrt{m^2 - m_0^2}$

**b**  $v = \frac{\sqrt{8}}{3}c$  **c**  $2.998 \times 10^8$  m/s

**EXERCISE J**

**1 a**  $\frac{15+x}{5}$  **b**  $\frac{x+3}{x}$  **c**  $\frac{x+4}{2}$  **d**  $\frac{14-x}{4}$

**e**  $\frac{8x-2}{15}$  **f**  $\frac{4x+17}{12}$

**2 a**  $\frac{x+5}{x+2}$  **b**  $\frac{11-2x}{x-4}$  **c**  $\frac{1-3x}{x-1}$  **d**  $\frac{5x+2}{x+1}$

**e**  $\frac{2x+3}{x+1}$  **f**  $\frac{x+3}{1-x}$

**3 a**  $\frac{7x^2-6x-25}{(2x-5)(x-2)}$  **b**  $\frac{-1}{(x-2)(x-3)}$

**c**  $\frac{8x^2+6x+8}{x^2-16}$  **d**  $\frac{3x^2+3x+13}{(x-3)(2x+1)}$

**EXERCISE K.1**

**1 Hint:** Consider  $\triangle$ s AEC, BDC

**2 Hint:** Let M be on AB so that  $PM \perp AB$

Let N be on AC so that  $PN \perp AC$

Join PM, PN and consider the two triangles formed.

**3** No hint needed.

**EXERCISE K.2**

**1 a**  $x = 2.4$  **b**  $x = 2.8$  **c**  $x = 3\frac{3}{11}$  **d**  $x = 6\frac{2}{3}$   
**e**  $x = 7$  **f**  $x = 7.2$

**2** 1.35 m tall

**EXERCISE L**

**1 a**  $\sqrt{13}$  units **b**  $\sqrt{34}$  units **c**  $\sqrt{20}$  units **d**  $\sqrt{10}$  units

**2 a** (2, 3) **b** (2, -1) **c**  $(3\frac{1}{2}, 1\frac{1}{2})$  **d** (2,  $-2\frac{1}{2}$ )

**3 a**  $\frac{2}{3}$  **b**  $-\frac{2}{5}$  **c** 1 **d** 0 **e** -4 **f** undefined

**4 a** 2 **b** 3 **c** undefined **d** 0 **e** -4 **f**  $\frac{1}{4}$

**5 a** parallel, slopes  $\frac{2}{5}$  **b** parallel, slopes  $-\frac{1}{7}$

**c** perpendicular, slopes  $\frac{1}{2}, -2$  **d** neither, slopes -4,  $\frac{1}{3}$

**e** neither, slopes  $\frac{2}{7}, \frac{1}{5}$  **f** perpendicular, slopes 2,  $-\frac{1}{2}$

**6 a**  $-\frac{4}{3}$  **b**  $-\frac{3}{11}$  **c**  $-\frac{1}{4}$  **d** 3 **e**  $\frac{1}{5}$  **f** undefined

**7 a**  $y = 2x - 7$  **b**  $y = -2x + 4$  **c**  $y = 3x - 15$

**d**  $y = -3x + 4$  **e**  $y = -4x + 9$  **f**  $y = x + 5$

**8 a**  $3x - 2y = 4$  **b**  $3x + 2y = 11$  **c**  $x - 3y = 4$

**d**  $4x + y = 6$  **e**  $3x - y = 0$  **f**  $4x + 9y = -2$

**9 a**  $x - 3y = -3$  **b**  $5x - y = 1$  **c**  $x - y = 3$

**d**  $4x - 5y = 10$  **e**  $x - 2y = -1$  **f**  $2x + 3y = -5$

**10 a**  $y = -2$  **b**  $x = 6$  **c**  $x = -3$

	Equation of line	Slope	x-int.	y-int.
<b>a</b>	$2x - 3y = 6$	$\frac{2}{3}$	3	-2
<b>b</b>	$4x + 5y = 20$	$-\frac{4}{5}$	5	4
<b>c</b>	$y = -2x + 5$	-2	$\frac{5}{2}$	5
<b>d</b>	$x = 8$	undef.	8	no y-int.
<b>e</b>	$y = 5$	0	no x-int.	5
<b>f</b>	$x + y = 11$	-1	11	11
<b>g</b>	$4x + y = 8$	-4	2	8
<b>h</b>	$x - 3y = 12$	$\frac{1}{3}$	12	-4

**12 a** yes **b** no **c** yes

**13 a** (4, 2) **b** (-2, 3) **c** (-3, 6) **d** (4, 0)

**e** parallel lines do not meet **f** coincident lines

**14 a**  $y = -4$  **b**  $x = 5$  **c**  $x = -1$  **d**  $y = 2$

**e**  $y = 0$  **f**  $x = 0$

**15 a**  $3x - 4y = -19$  **b**  $x - y = 7$  **c**  $y = 3x$

**d**  $2x + 3y = 19$  **e**  $x - 2y = 5$  **f**  $x + 3y = 13$

**16 a**  $x - 8y = -83$  **b**  $8x + y = 41$

**c**  $9x - 2y = 23$  for  $5 \leq x \leq 7$  **d**  $(8, 11\frac{3}{8})$

**17 a**  $x - 3y = -16$  **b**  $2x - y = -3$  **c**  $x = 5$

**18**  $(4\frac{3}{4}, 8)$

**19 a**  $(2\frac{1}{3}, 7)$  **b** 8.03 km **c** yes (6.16 km)

**20** (13.41, 8) or (-5.41, 8)

**21 Hint:** Use the distance formula to find the distance from the centre of the circle to point P.

**22 a**  $(x-4)^2 + (y-3)^2 = 25$  **b**  $(x+1)^2 + (y-5)^2 = 4$

**c**  $x^2 + y^2 = 100$  **d**  $(x-1)^2 + (y-3)^2 = 8$



- 23** **a** centre (1, 3), radius 2 units  
**b** centre (0, -2), radius 4 units  
**c** centre (0, 0), radius  $\sqrt{7}$  units
- 24** **a** centre (2, -3), radius  $\sqrt{20}$  units  
**b** **Hint:** Substitute (4, 1) into equation of circle.  
**c**  $x + 2y = 6$
- 25** (3, 3)